



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Embryo of *Helminthostachys*.—LANG³⁴ has supplied some much needed information in reference to the embryogeny of *Helminthostachys*. In a preliminary note³⁵ he had announced the existence of a well developed suspensor, but the present study furnishes many additional details. The embryo extends down into the prothallium before segmentation takes place, and the first two walls are transverse. The cell next to the neck of the archegonium, which may divide or not, forms the upper suspensor tier; the middle cell, which divides, forms the second suspensor tier; while the terminal cell forms the embryo proper. The hypobasal half of the embryo forms the foot, while from the epibasal half the stem tip, first leaf, and probably the first root arise. A comparative study of the embryogeny of Marattiaceae, Ophioglossaceae, and seed plants leads to the suggestion that "the suspensor represents the last trace of the filamentous juvenile state in development of the plant, and may have persisted in the seed-plants from their filicineous ancestry."—J. M. C.

Basidiomycetes of the Philippines.—GRAFF³⁶ has published a list of additions to the known Basidiomycetes of the Philippines with descriptions of new species. These additions number 33, about equally distributed between Hymenomycetes and Gasteromycetes. The new species are described in *Exidia*, *Laschia*, *Lentinus*, *Volvaria*, *Naucoria*, and *Bovista*.—J. M. C.

Nectaries and phylogeny.—After examining the nectaries of a large number of monocotyledons and dicotyledons, PORSCH³⁷ reaches the conclusion that the nectary is not only an organ of some phylogenetic significance, but that it furnishes additional proof of the derivation of the former from the latter.—C. J. CHAMBERLAIN.

³⁴ LANG, WILLIAM H., Studies in the morphology and anatomy of the Ophioglossaceae. II. On the embryo of *Helminthostachys*. Ann. Botany 28:19-37. figs. 9. pl. 3. 1914.

³⁵ Ann. Botany 24:611. 1910.

³⁶ GRAFF, P. W., Additions to the basidiomycetous flora of the Philippines. Philippine Jour. Sci. 8:299-307. pls. 8-10. 1913.

³⁷ PORSCH, OTTO, Die Abstammung der Monokotylen und die Blütennektarien. Ber. Deutsch. Bot. Gesells. 31:580-590. 1914.